ISSN 0013-8908

THE ENTOMOLOGIST'S MONTHLY MAGAZINE

World List Abbreviation, 4th ed. (1963): *Entomologist's mon. Mag.*

EDITED BY
Prof. Sir Richard SOUTHWOOD, D.Sc., F.R.S., F.R.E.S.
Prof. M. G. MORRIS, Ph.D., F.I.Biol., F.L.S., F.R.E.S.
K. G. V. SMITH, C.Biol., F.I.Biol., F.L.S., F.R.E.S.
J. W. ISMAY, B.Sc., Ph.D., F.R.E.S.
P. M. HAMMOND, M.A., F.R.E.S.
W. R. DOLLING, B.Sc., F.R.E.S.

VOLUME 134

'J'engage donc tous à éviter dans leurs écrits toute personnalité, toute allusion dépassant les limites de la discussion la plus sincère et la plus courtoise.' — *Laboulbène.*

THE ENTOMOLOGIST'S MONTHLY MAGAZINE
BRIGHTWOOD
BRIGHTWELL, WALLINGFORD, OXON OX10 0QD
1998

Published by Gem Publishing Company

CONTENTS

CONTRIBUTORS iii
GENERAL INDEX iv
SPECIAL INDEX –
ARACHNIDA vii
COLEOPTERA vii
COLLEMBOLA viii
DERMAPTERA viii
DIPTERA viii
EPHEMEROPTERA xiii
HEMIPTERA xiii
HYMENOPTERA xiii
ISOPTERA xv
LEPIDOPTERA xv
MECOPTERA xv
ODONATA xv
ORTHOPTERA xv
PSOCOPTERA xv
THYSANOPTERA xvi
TRICHOPTERA xvi
GROUPS OTHER THAN INSECTA AND ARACHNIDA xvi
ERRATA AND CORRIGENDA xvi

DATES OF PUBLICATION

January/February/March/April	27th March 1998
May/June	30th June 1998
July/August	31st July 1998
September/October/November/December	30th November 1998

INDEX

This index covers the year 1998 (Vol. **134**). Items on the wrappers are indicated by Roman numerals. Prepared by Mrs A.V. Smith.

INDEX TO CONTRIBUTORS

Aldridge, M.C., 292
Alexander, K.N.A., 9
Allen, A.A., 213, 214, 270
Angelini, F., 329
Angus, R.B., 5
Archer, M.E., 38, 45, 215, 238, 263, 279, 306, 334
Ashe, P., 338
Askew, R.R., 53
Baldwin, A.J., 325
Beaver, R.A., 285
Blackburn, J.H., 94, 257
Blackman, R., 44
Booth, R.G., 335
Bowden, J., 61
Buhl, P.N., 241
Carter, D., 310
Cole, J., 84
Collins, D.W., 1
Colston, A., 353
Cooper, M., 89, 293
Cooter, J., 329
Corbet, S.A., 4, 65, 344
Deeming, J.C., 44, 314
Denton, J.S., 265, 327, 350
Dillon, R., 229
Disney, R.H.L., 231, 344, 356
Donnelly, M.J., 65
Eds., xvii
Foster, G.N., 264
Fry, R.M., 327
Goldarazena, A., 319
Griffiths, G.C.D., 328
Gunn, R.J.M., 94, 257
Hammett, M.J., 257
Hammond, P.M., 273
Hancock, E.G., 223
Harding, P.T., 230, 264
Holloway, J., 256
Hopkins, G.W., 239, 271
Horsfield, D., 352
Ismay, J.W., 344, 351
Judd, S., 311
Kassebeer, C., 39
Lampe, K.-H., 53
Laurence, B.R., 264, 334
McLean, I.F.G., 85
McManus, C.E.L., 75
Maibach, A., 39
Maier, C.T., 223
Marullo, R., 345
Miles, P.M., 11, 334
Morris, M.G., 197, 277
Mound, L.A., 319
Nagatomi, A., 95
Nash, R., 4, 255
Nau, B.S., 52
Nelson, J.M., 77
O'Connor, J.P., 4, 93, 255, 284, 338
O'Connor, M.A., 284
Palmer, J.L., 65
Paul, J., 60
Peacey, M., 351
Plant, C.W., 270
Pont, A.C., 193, 344
Rotheray, G.E., 39, 223
Saini, M.S., 69, 315
Shaw, M.R., 52
Simmonds, M., 228
Smith, K.G.V., 85, 88, 238, 358
Smithers, C.N., 57
Southwood, T.R.E., 194, 357
Staddon, B.W., 267
Sullivan, C., 30
Valladares, G.R., 31, 235
Vasu, V., 69, 315
Verdcourt, B., 43
Webb, N.R., 355
Welch, R.C., 272, 339
Whitehead, P.F., 30, 214, 266, 307
Whittington, A.E., 278
Wistow, S., 338
Wood-Baker, C.S., 239, 271
Wright, P.J., 75, 354
Yang, D., 95

GENERAL INDEX

A.A. Allen's 85th birthday, xvii
Aerial distribution of puparium of *Leucopis glyphinivora* Tan. (Dipt., Chamaemyiidae) via thistle pappus, 85
Antennal sensoria, diversity of, amongst species in the basal clades of the Thysanoptera, 345
Antennal teratology (Col.) again, 214
Arabia, The tropical nest fly, *Passeromyia heterochaeta* (Villeneuve) (Dipt., Muscidae) in eastern, 314
Argentina, a new species of *Amauromyza* Hendel (Dipt., Agromyzidae) from, 235; three new species of *Haplopeodes* Steyskal (Dipt., Agromyzidae) from, 31
Aspen, *Populus tremula* L., comparative aspects of the biology of three species of *Dorytomus* (Col., Curculionidae) associated with, 197
Australia, a year's Malaise trap catches of Psocoptera in, 57
Bedfordshire, *Formica cunicularia* Latr. in, an update, 354; further notes on range-changes of Hemiptera-Heteroptera in, 52; root aphids and ants in grassland, 75
Berkshire, *Phasia hemiptera* (F.) (Dipt., Tachinidae) at Maidenhead, 43
Britain, An Asian rove-beetle, *Oxytelus migrator* Fauvel (Col., Staphylinidae) established in, 273; *Caenis beskidensis* Sowa (Ephem., Caenidae), a mayfly new to, 94; *Cyphea curtula* (Erichson) (Col., Staphylinidae) new to, 339; *Electrogena affinis* (Eaton 1885) (Ephem., Heptogenidae) a mayfly new to, 257; four species of Agromyzidae new to, 84; *Leucopis glyphinivora* Tanasijtshuk (Dipt., Chamaemyiidae) new to, and the aerial distribution of its puparium via thistle pappus, 85; *Neorhegmoclemina catharinae* Haenni (Dipt., Scatopsidae) new to, 351
British records of *Eurysa brunnea* Melichar, 1896 (Hem., Delphacidae), 307
British seed bugs (Hem., Lygaeidae), new national, regional and county records of, 311
Cambridgeshire and Huntingdonshire, the range expansion of *Metrioptera roeselii* Hagenbach (Orth, Tettigonidae) in, 353
Cardiganshire (Wales) the Collembola of, 11; *Labia minor* L. (Derm., Labiidae) from the summit of Yr Wyddfa (Mt Snowdon), 30
Cornwall, rare and scarce wasps and bees (Hym., Aculeata) found in during 1996 and a plea for help, 38
Cotswold Hills, a national stronghold for *Stenostola dubia* (Laich.) (Col., Cerambycidae), 9
Devon, *Ptenidium brenskei* Flach (Col., Ptilidae) in south, 213
Dipterist's Forum, xii
Drought, behaviour of *Episyrphus balteatus* (De Geer) (Dipt., Syrphidae) during, 334
England, rare and notable Hemiptera in southern, 1995–97, 265; recent interceptions of *Echinothrips americanus* Morgan (Thys., Thripidae) imported into, 1; recent records of scarce and notable hoverflies (Dipt., Syrphidae), 327
Essex and nearby, more on *Didea fasciata* Macquart (Dipt., Syrphidae) in, 270
Euro-Asian distribution of species of *Paravespula* (Hym., Vespidae), 279
Extinct Mesozoic genera and families of Brachycera (Insecta, Diptera, Orthorrhapha), a review of, 95
Fecundity and development, effect of magnetic field on, in *Subcoccinella vigintiquattuorpunctata* (L.) (Col., Coccinellidae), 325
Feeding on mosquitoes, *Hydrometra stagnorum* (L.) (Hem., Hydrometridae), 65
Frequency of *Lagria hirta* (L.) (Col., Lagriidae), have you noticed exceptional, recently, 264
Galls of Cecidomyiidae (Dipt.), species of Pteromalidae (Hym., Chalcidoidea) parasitic in, on *Fagus* leaves, 53
Guernsey and Jersey, misidentifications of solitary bees (Hym., Aculeata) from, 334
Gynandromorph of *Halictus confusus* Smith (Hym., Halictidae), 238
Hampshire, recent records of scarcer Aculeate Hymenoptera and Surrey and North, 350
Hertfordshire, *Volucella zonaria* (Poda) (Dipt., Syrphidae), a recent extension of its range into, 292

Huntingdonshire and Cambridgeshire, the range expansion of *Metrioptera roeselii* Hagenbach (Orth., Tettigonidae) in, 353
India, a new genus of Blennocampinae (Hym., Symphyta, Tenthredinidae) from, 69
Ireland, *Anomoia purmunda* (Harris) (Dipt., Tephritidae) swarming in a freshly painted shed, 284; *Dermestes peruvianus* Laporte de Castelnau (Col., Dermestidae) new to, 93; *Drosophila busckii* Coquillett (Dipt., Drosophilidae) an indoor record in, 338; *Pygostolus multiarticulatus* (Ratzeburg) (Hym., Braconidae) new to, 255
Irish records, further, of *Diastrophus rubi* (Bouché) (Hym., Cynipidae), 4
Isle of Wight, *Chrysolina viridana* (Küster) (Col., Chrysomelidae) found in, 270
Jersey and Guernsey, misidentifications of solitary bees (Hym., Aculeata) from, 334
Larva (third = final), stage of *Psilota anthracina* Mg. and *Psilota decessa* (Hutton) (Dipt., Syrphidae), 39
London (S.E.) garden, *Didea fasciata* Macq. (Dipt., Syrphidae) in a, 270
Magnetic field, effect of, on fecundity and development of *Subcoccinella vigintiquattuor-punctata* (L.) (Col., Coccinellidae), 325
Malaise trap catches of Psocoptera, a year's, in Australia, 57
Mesozoic (extinct) genera and families of Brachycera (Insecta, Diptera, Orthorrhapha), a review of, 95
Middlesex, a recent record of *Meroplius minutus* (Wiedemann) (Dipt., Sepsidae) from, 344
New species of the *artifex* group of *Mischocyttarus* de Saussure (Hym., Vespidae) with a partial key, 293; of *Mischocyttarus* (Hym., Vespidae) with notes on some members of the *iheringi* group, 89
Norfolk Broads, *Hydroporus glabriusculus* Aubé (Col., Dytiscidae) new for the, 264
Nottingham, *Rhynchium oculatum* (F.) (Hym., Eumenidae) found at, 263
OBITUARY – Dr Derek A. Whiteley, 88
Observations of *Neurigona abdominalis* (Fallén) and other Dolichopodidae (Dipt.), 61
Oriental *Spathidicerus* Chapuis (Col., Platypodidae) three new, with a key to species, 285
Oviparous female of *Laingia psammae* Theobald 1922 (Hem., Aphididae), 271
Painted Irish shed, *Anomoia purmunda* (Harris) (Dipt., Tephritidae) in a freshly, 284
Perth and Mull, Scotland, *Botanophila biciliaris* (Pandellé) (Dipt., Anthomyiidae) from the Tay reed beds, 352
REVIEWS — **Abrahamson, W.G. & Weis, A.F.**, 1997, Evolutionary Ecology across three trophic levels, 194; **Acta Entomologica Serbica** (1997), **2**, 324; **Andersen, S.**, 1996, The Siphonini (Diptera: Tachinidae) of Europe (*Faun. ent. Scand.* **35**), 238; **Annual Reviews of Ecology and Systematics**, 1997, **28**, xii; **Butterfly Conservation in Hertfordshire & Middlesex**, 1997, 310; **Day, K.R. et al.**, 1998, The Green Spruce Aphid in Western Europe (Forestry Commission), 318; **De Vries, P.J.**, 1997, The Butterflies of Costa Rica and their Natural History, Vol. 2, Riodinidae, 256; **Diedrich, J.L.T.**, 1997, Contribution à la Connaissance des Cétoines (Col., Cetoniinae) de la Sous-Region Malgache, xvii; **Edwards, R.** (ed.), Provisional Atlas of the Aculeata Hymenoptera of Britain and Ireland, part 1 (1997), v; part 2 (1998), xxi; **Evaluation of Genetically altered medflies for Sterile Insect Technique Programmes**, 1997, (International Atomic Energy Agency), 344; **Entomotaxonomia**, 1997, **19**(3–4), 192; **Evenhuis, N.L.**, 1997, Litterature Taxonomica Dipterorum (1758–1930), 193; **Food Insects Newsletter** 1997, **10**(1), 51; **Gaedike, R.**, 1996, Lepidopterorum Catalogus (New Series) **4**(48) Epermenidae, 256; **Gangwere, S.K., Muralirangan, M.C. & Muralirangan, M.**, 1997, The Bionomics of Grasshoppers, Katydids and their Kin, 355; **Haes, E.C.M. & Harding, P.T.**, 1997, Atlas of grasshoppers, crickets and allied insects in Britain and Ireland, 60; **Hansen, M.**, 1997, Phylogeny and Classification of the Staphyliniform beetle families (Coleoptera) (*Biol. Skr. Dan. Vid. Selsk; Biologiske Skrifter*, **48**), 272; **Heinrich, B.**, 1996, The Thermal Warriors, 4; **I.B.R.A.**, 1997, Varroa! Flight the mite, 3; **INRA – ACTA** 1998, Les Pucerons des Arbres Fruitiers, 349; **Jolivet, P.H.A. & Cox, M.L.** (eds), 1996, Chrysomelid Biology, 277; **Karban, R. & Baldwin, I.T.**, 1997, Induced Responses to Herbivory, 194; **Lehane, M.J. & Billingsley, P.F.** (eds), 1996, Biology of the Insect Midgut, 228; **Mason, F.**, 1997, The Afrotropical Nemotelinae (Dipt., Stratiomyidae), *Monografia* **XXIV**, Museo Regionale de Scienze

Naturali, Torino, 328; **Morris, R.K.A.**, 1998, Hoverflies of Surrey, 354; **Myia**, 1979–1994, Summary of Contents, xi; **National Geographic Society**, 1997, Last Wild Places, 357; **Nijhout, H.F.**, 1997, Insect Hormones (pbk edn), xvii; **Nilsson, A.**, 1997, Aquatic Insects of North Europe, **2** (Odonata – Diptera), 356; **Oistros**, 1997, Summary of Contents, xi; **Porter, J.**, 1997, The Colour Identification Guide to Caterpillars of the British Isles (Macrolepidoptera), 358; **Price, J.M.**, 1997, A Provisional Atlas of the True Bugs of Warwickshire (Hem. Het.), xvii; **Remaudière, G. & M.**, 1997, Catalogue des Aphididae du Monde (I.N.R.A.), 44; **Rothschild, M. & Marren, P.**, 1997, Rothschild's Reserves, 357; **Smith, F.H.N.**, 1997, The Moths and Butterflies of Cornwall and the Isles of Scilly, 310; **Stadler, E., Rowell-Rathier, M. & Baur, R.** (eds), 1996, Proceedings of the 9th International Symposium on Insect-Plant Relationships (*Series Entomologica*, **53**), 194; **Stork, N.E., Adis, J. & Didham, R.K.** (eds), 1997, Canopy Arthropods, 194; **Thorens, P. & Nadig, A.**, 1997, Atlas de Distribution des Orthoptères de Suisse (*Documenta Faunistica Helvetiae*, **16**), 230; **Volucella**, 1995-, xi; **Waldbauer, G.**, 1998, Insects through the Seasons, pbk edn, 196; **Watt, A.D., Stork, N.E. & Hunter, M.D.** (eds), 1997, Forests and Insects, 194; **Wikel, S.** (ed.), 1996, The Immunology of Host-parasitic Arthropod relationships, 229

Scotland, *Botanophila biciliaris* (Pandellè) (Dipt., Anthomyiidae) from the Tay reed beds, Perth and Mull, 352; *Thanatophilus dispar* Herbst (Col., Silphidae) found on Uist, 30; two species of *Omalus* (Hym., Chrysididae) in, 52

Shetland, correction: *Pegomya argyrocephala* Mg. (Dipt., Anthomyiidae) not from, 264

Shropshire, Wyre Forest, extraordinary numbers of *Polyxenus lagurus* (L.) (Myriapoda) in the, 266

Spain, a new species of *Hindsiothrips* from, with the first record of *Tylothrips osborni* (Hinds) from Europe (Thysanoptera, Phleaeothripidae), 319

Static build-up in photo-canisters: a note of warning, 278

Surrey, recent records of scarcer aculeate Hymenoptera in North Hampshire and, 350

Taxonomic status of *Longitarsus bearei* Kevan (Col., Chrysomelidae), 335

Taxonomic studies on *Tomostethus gracilicornis* (Rohwer) (Hym., Tenthredinidae, Blennocampinae), 315

Taxonomy, distribution and nesting biology of *Vespa analis* F. (Hym., Vespidae), 215; of *Vespa orientalis* L. (Hym., Vespidae), 45

Thailand, descriptions of two new species of Agathidiini Westwood 1838 (Col., Leiodidae) new to, 329

Tree sap, the larvae of two *Ceriana* species (Dipt., Syrphidae) breeding in, 223

Turkey, colour and sternal patch pattern variation in a spring aggregation of *Nezara viridula* L. (Hem., Het., Pentatomidae) from, 267

Turkish *Helophorus*, a new, with notes on *H. griseus* Hbst and *H. montanus* d'Orch. (Col., Hydrophiloidea), 5

Uist [Scotland], *Thanatophilus dispar* Herbst (Col., Silphidae) found on, 30

U.S.A., two new species of *Puliciphora* Dahl (Dipt., Phoridae) from the, and an Old World synonym, 231

Wales, Collembola of Cardiganshire, 11; *Labia minor* (Derm., Labiidae) from the summit of Yr Wyddfa [Mt Snowdon], 30; the mosquito *Aedes (Finlaya) geniculatus* (Olivier) in South, 44

Welsh coastal towns, immigrant *Episyrphus balteatus* (De Geer) (Dipt., Syrphidae) and insects in, 334

Whale, Diptera trapped in the oil oozing from a dead, 344

Wiltshire, a record of the hornet (*Vespa crabro* L., Hym., Vespidae) from, with a plea for help, 306

Worcestershire, *Tribolium castaneum* (Herbst) (Col., Tenebrionidae) infesting dried insects in, 214

World distribution of Euro-Asian species of *Paravespula* (Hym., Vespinae), 279

World species of *Euxestonotus* Fouts (Hym., Platygastridae), key to, 241

SPECIAL INDEX

New genera and species, etc., are indicated by heavy type; genera and species new to Britain by an asterisk (*).

ARACHNIDA

Eriophyidae, 192
Eriophyoidea, 324
Ixodidae, 324
Varroa, 3

COLEOPTERA

Abax parallelipipedus, 214
'Aclypea opaca', 30
Adephaga, 272
Afroagathidium, 330
Agathidium argutum, 333; darbyi v. darbyi, 332–3; insolitum, laticorne, 333; kuaiense, 329, 331–2; kualalumpense, luctuosum, madurense group, 332; nigripenne, 343; schwendingeri, watrousi, wheeleri, 332
Amphimallon solstitiale, 214
Anomognathus, 340
Anotylus glareosus, mimulus, nitidifrons, pygmaeus, vinsoni, 275
apateticids, 272
Aphodius holdereri, 8; lapponum, 30
Archostemmata, 272
Aridius bifasciatus, nodifer, 343
Atheta nigra, 343; tibialis, 30
Attagenus pellio, 93
Bembidion, 214
Bostrichiformia, 272
Carabus violaceus, 214
Cetoniinae, xvii
Chrysolina *viridana, 270
Chrysomelidae, 277–8
Coccinella 7-punctata, 334
Cryptolestes ferrugineus, 343
Cryptophagus dentatus, intermedius, 343
Cupepidae, 272
Curculionidae, 255, 324
Cyphea *curtula, 339–43; latiuscula, 339–40
Dasycerinae, 272
Datomicra, 343
Dermestes peruvianus, 93
Derodontidae, 272
Dorytomus affinis, 197–8, 200–1; dejeani, 197–9, 201–5, 207–12; edoughensis, 197; luridus, 197; tortrix, 197–9; tremulae, 197, 211
Dropephylla ioptera, 343
dung beetles, 4
Elateriformia, 272
empelids, 272
Enicmus brevicornis, 339, 343
Etiophilus, 197
Epimetopidae, 272
Eucinetidae, 272
Geodromicus longipes, 30
Georissidae, 272
Gyrinidae, 272
Gyrophaena, 342
Haplogastra, 272
Harpalus rubripes, 308
Helophoridae, 272
Helophorus, 272; aquaticus, aspericollis, browni, 8; carsoni, 7; discrepans, glacialis, granularis, 8; griseus, helenae, 5–8; jacatus, 8; montanus, 5–8; paramontanus, 8
Histeroidea, 272
Homalota, 340; plana, 343
Hydraenidae, 272
Hydrochidae, 272
Hydrophilidae, Hydrophiloidea, 272
Hydroporus glabriusculus, scalesianus, 264
Laccornis oblongus, 264
Lagria hirta, 264
Leiodidae, 272
Leperesinus orni, varius, 343
Leptinus, 272
Leptusa fumida, norvegica, ruficollis, 343
Longitarsus agilis, ballotae, 336; bearei, 335–7; lewisii, luridus, medvedevi, 337; plantago-maritimus, 336; pratensis, 335–7; pratensis group, 335–6
Mecopelus, 291
Microceble, 333
Micropeplinae, 272
Microsporidae, 272
Myxophaga, 272
Nebria gyllenhali, 30
Ommatidae, 272
Otiorhynchus singularis, 255
Oxytelus akazawensis, 273; alutaceifrons, depauperatus, 275; *migrator, 273–5; piceus, 276; puncticeps, sculptus, 275
Patrobus assimilis, 30
Perigona nigriceps, 275
Periommatus, 285
Phloeonomus punctipennis, 343
Phytoecia cylindrica, 10
Placusa, 340
Platypsyllinae, 272
Pselaphidae, 272

Pseudoagathidium burgeoni, congolense, 329; **immigrans**, 329–30
Ptenidium brenskei, intermedium, 213
Ptiliidae, 272
Rhizophagus bipustulatus, 343
Saigonium quadricorne, 339, 342–3
Saperda perforata, 343
scaphidiids, 272
Scarabaeoidea, 272
Schedlarius, 291
scydmaenids, 272
Spathidicerus, 285–91; **brownei**, 286–90; corporaali, 286, 288–9; intermedius, 285; javanus, 286–7, 291; nobilis, 285–6; papuanus, philippinensis, 286–7; **sulawesianus**, 286–7, 289–90; sumatranus, 286–9, 291; **suttoni**, 286–7, 290–1; thomsoni, 285–7
Sperchidae, 272
Staphyliniformia, 272
Staphylinoidea, 272
Stenostola dubia, 9–10
Subcoccinella vigintiquattuorpunctata, 325–7
Tachyusa, 342
Tesserocerus, 285
Thanatophilus dispar, 30
Thectura cuspidata, 342
Trachypachidae, 272
Tribolium castaneum, 214
Trogidae, 272
Trypophloeus, 343
Typhaea stercorea, 30
Xantholinus, 214
Xyloterus, 343

COLLEMBOLA

Allacma, 29
Anurida granaria, 19
Anurophorus laricis, unguiculus, 21
Arrhopalites principalis, 29
Bourletiella pallipes, 29
Brachystomella parvula, 18–9
Ceratophysella armata, bengtssoni, denticulata, longispina, 17
Collembola, 192, 324
Deuterosminthurus, 29
Dicyrtomina minuta, 29
Entomobrya albocincta, lanuginosa, 25; multifasciata, 25–6; nivalis, 26
Folsomia bisetosa, brevicauda, 21; diplophthalma, 21–2; fimetaria, fimetarioides, penicula, quadrioculata, sexoculata, 22
Friesea claviseta, mirabilis, 17
Heteromerus nitidus, 27
Hypogastrura purpurescens, sahlbergi, viatica, 17
Isotomiella minor, 22
Isotomina arborea, 25; maritima, notabilis, 23; olivacea, 23–4; o.f. tigrina, 24, 24; sensibilis, 24–5; thermophila, 23; violacea, viridis, 24
Isotomodes productus, 21
Isotomurus palustris, 25
Kalophorura burmeisteri, 20
Lathriopyga grassei, 19
Lepidocyrtus curvicollis, cyaneus, 26; lanuginosus, 26–7
Megalothorax, 28
Mesophorura, 20
Metaphorura, 20
Neanura muscorum, 19
Neelus minimus, murinus, 28
Odontella *empodialis, 17
Onychiurus ambulans, fimetaroides, 19
Orchesella alticola, cincta, villosa, 27
Paratullbergia, 20
Podura aquatica, 17
Pogonognathellus, 28
Protophorura armatus, 19–20; fimatus, 20
Pseudachorutes corticicolus, dubius, subcrassus, 19
Pseudisotoma, 24–5
Schaefferia willemi, 17–8
Schoettella ununguiculata, 17
Seira domestica, 27
Sminthurides malmgreni, parvulus, 28; pumilis, 28–9; violaceus, 29
Sminthurinus aureus, elegans, 29
Sminthurus fuscus, viridis, 29
Stenacidia, 29
Tetracanthella pilosa, 20; wahlgreni, 20–1
Tomocerus longicornis, 28; minor, 27; minutus, 27–8; vulgaris, 27–8
Tullbergia affinis, callipygos, concolor, krausbaueri, quadrispina, 20
Vertagopus, 25
Xenylla brevicauda, corticalis, humicola, maritima, 18
Xenyllodes armatus, 18

DERMAPTERA

Labia minor, 30

DIPTERA

Acroceridae, 95–6, 156, 159, 162–3, 172, 187–90, 192
Actia nudibasis, resellae, 238
Aedes aegypti, 324; geniculatus, 44
Agromyza albipennis, *graminicola,

intermittens, *luteitarsis, 84
Akronia, 328
Alinka, 101; cara, 103
Alinkidae, 101, 103–4, 148, 158
Alleremonomus, 173–4, 177; liaoningensis, 174; xingi, 173–4, 187
Alloleptis, 98
Alophora (s.g.), 43
Amauromyza fuscibasis, 235; maculosa, 235, 237; **pterocaula**, 236–7
Anacanthaspis, 99
Anisopodidae, 104, 160, 162
Annimyzella (s.g.), 235, 237
Anomoia purmunda, 284
Anthomyiidae, 193, 344
Antissini, 118
Apioceridae, 179, 186
Apsilocephala, 118–20; longistyla, 149–50, 153
Apsilocephalidae, 95–7, 116, 118, 120, 154, 156
Apystomyiidae, 96
Archinemestriinae, 180, 183
Archinemestrius, 182–3, 189; karatavicus, 179–80, 184, 187
Archiphora ancestrix, 115
Archiphoridae, 115
Archirhagio, 101, 105, 110, 156, 164; obscurus, 105–6, 171, 187
Archisargidae, 100–1, 103, 105–9, 154, 156, 158, 162–71, 187–90
Archisargoidea, 163
Archisargus, 103, 105, 156, 164, 166–7, 171; maximus, 164–5; pulcher, 106–7, 164–5
Archisolva cupressa, 139–40
Archocyrtidae, 162–3, 172, 185, 187–90
Archocyrtinae, 172
Archocyrtus, 162; gibbosus, 172, 187–8, 192
Argyra argentella, diaphana, leucocephala, 64
Arthroceras, 99, 135
Arthropeas, 99
Arthroteles 99, 135
Aschiza, 174
Asilidae, 95–6, 156, 159
Asiloidea, 95–6, 156, 158, 186
Asilomorpha, 192
Athericidae, 96, 99–100, 133, 142, 147, 356
Atherimorpha, 99, 135
Atriadopsinae, 188
Atrichops, 99
Austroleptis, 98, 136
Batgana, 102, 141–2, 158; minor, 141–2
Bequaertomyia, 185
Bibionidae, 104, 192
Bolbomyia, 98, 119, 122, 127–9, 132–3, 136
Bombyliidae, 95–6, 110–1, 136, 156, 158–9
Botanophila biciliaris, 352
Brachycara, 328
Brachycera, 95–192
Brachyopa, 226; insensilis, 327, pilosa, testacea, 42
Brachypalpoides lenta, 42
Brachypalpus, 226
Butomyza (s.g.), 84
Calliphora vicina, xi
Calliphoridae, 156, xi
Calosargus, 164; niger, 165–6, 170; tatianae, 165–6, 170, 187; thanasymus, 165–6, 170
Campsicnemus curvipes, 64
Camposella, 188
Camptopelta, 328
Cerantha, 238
Ceratinostoma ostiorum, 77–83
Ceratitis capitata, 344
Ceratopogonidae, 356
Ceriana abbreviata, 223–4, 226; eumenoides, 277; odontomera, 223–7
Cerodontha *scutellaris, 84
Chalcosyrphus, 226
Chaoboridae, 356
Chionea, 104
Chiromyzinae, 118
Chironomidae, 344, 356
Chyromya flava, oppidana, 338
Chrysogaster virescens, 327
Chrysopilus, 100, 134, 136
Chrysotoxum octomaculatum, 327
Chrysotus blepharosceles, 64; varians, 62, 64
Clesthentia, 119–20; sp., 151–2, 155
Clesthentiella, 119–20, 157; crassioccipitis, 153, 157
Clitellariinae, 118, 328
Cluninemotelus, 328
Coelopa, 79; frigida, 344
Coenomyia, 99
Coenomyiidae, 96, 99, 127–8, 133, 159, 186
Cramptonomyiidae, 104
craneflies, 356
Cretaceogaster, 101, 138–9, 161; pygmaeus, 138–9
Criorhina asilica, ranunculi, 327
Culex pipiens f. molestus, 65–8
Culicidae, 356
Cyclopsideinae, 188
Cyclorrhapha, 95–7, 156, 158

Dasyomma, 100
Dialysis, 99, 127, 133, 159
Didea fasciata, 270
Diptera, 193
Dissup, 105, 111–2, 173, 177
Dixidae, 356
Dizygomyza (s.g.), 84
Dolichopodidae, 96, 155, 159, 356
Dolichopus festivus, latelimbatus, 62; ulgulatus, 64; wahlbergi, 62
Drosophila busckii, 338 melanogaster, 65; subobscura, 324
Drosophilidae, 324
Elachisoma aterrimum, 344
Empidia, 95
Empididae, 95–6, 117, 155–6, 159, 356, xi
Empidoidea, 95–7, 156, 158
Eohirmoneura, 189; carpenteri, 180–1, 183, 187
Eomyia, 102, 109; veterrima, 109–10
Eomyiidae, 102, 109–10, 152, 158
Eoptychopteridae, 123
Eoptychopterina daiensis, 123
Eostratiomyia, 102, 110; avia, 110
Eostratiomyiidae, 102, 110, 158
Ephydridae, 344, 356
Epideicticus, 328
Episyrphus balteatus, 334
Eremochaetidae, 100, 105, 111–5, 154, 156, 158, 162–3, 172–7, 185, 187–90
Eremochaetinae, 177
Eremochaetomima, 177; incompleta, 175
Eremochaetosoma, 105, 111, 113–4, 154, 177; mongolicum, 113, 187
Eremochaetus, 111, 113–4, 173, 177; asilicus, 113–4
Eremocreta (s.g.), 176–7, 187
Eremomukha, 177; addita, 176; insidiosa, 175; posita, sorosi, 176, 187; tsokotukha, 175–6, 187
Eremomukhinae, 177
Eremonomus irae, 112
Eumerus, 41
Eurosta solidaginis, 194
Exeretonevra, 179, 185–6
Exeretonevridae, 96, 98, 185–6
Exodontha, 118
Fallenia, 179
Ferdinandea, 226; ruficornis, 328
Finlaya (s.g.), 44
Fucellia, 79
Geitonomyia, 328
Gen. et sp. 1 [Kovalev], 95, 125–6; 156
Gen. et sp. 2 [Kovalev], 95, 125–6, 137
Glutops, 99–100, 110, 127–9, 133, 145, 185
Hammerschmidtia, 226
Hartigiola annulipes, 53–6
Haplopeodes capsici, 31–2; eurhabdus, 31; **flavinotus**, 31–3; **gomphrenae**, 31, 34–6; lopesi, 31; **lycivorus**, 31, 36–8; minutus, 36, 38; tigrensis, 31
Haplomyza, 31
Hasmona, 95
Hercostomus chalybeus, cupreus, germanus, metallicus, 64; ungulatus, 62–3
Heterostomidae, 96, 98, 186
Heterostomus, 186
Hilara, xi
Hilarimorphidae, 96
Hirmoneura, 180–3; willistoni, 180, 182
Hirmoneurinae, 181, 188
hoverflies, xi
Hypophyllus obscurellus, 64
Ija, 101–2, 127, 141–3, 152, 156, 158; problematica, 142–3
Inopus, 118
Ironomyiidae, 156
Jurabrachyceron, 102, 127–8, 137; abbreviatum, 127–8
Juracyrtus kovalevi, 192
Kerosargus argus, 178, 187–8
Kovalevisargidae, 162–3, 178–9, 187–9
Kovalevisargus clarigenus, 178–9
Kubekovia, 102, 127–9, 137; accessoria, 128–9
Lasiopa, 328
leatherjackets, 238
Lepidomyia decessa, 39–41
Leptocera coenosa, 344
Leptoynoma, 98
Leucopis *glyphinivora, 85–7; melanopus, puncticornis, 85
Liriomyza, 31; huidobrensis, 237
Litoleptis, 98
Lucilia, 334
Malacomyia sciomyzina, 344
Medetera flavipes, 64; jacula, 62, 64; muralis, petrophiloides, 64; truncorum, 62, 64
Megaselia diversa, nigriceps, pleuralis, 344
Melanostoma decessa, 39
Merodon, 41
Meroplius minutus, 344
Mesomphrale, 101, 141–3, 156, 158; asiaticum, 143–4
Mesorhagiophryne, 141, 144–5, 152, 156, 158; incerta, 102, 144; robusta, 102, 141, 145, 152
Mesosolva, 101, 105–6, 111, 154, 156, 163–4; angustocellulata, 166–7, 171; balyshevae, 166–7, 171, 187; dolosa, hennigi, imperfecta, longivena, 166–7,

171; parva, 107–8, 166, 169, 171, 187; rohdendorfi, 166–7, 171
Mesostratiomyia, 101, 141–2, 146, 152, 156, 158; laiyangensis, 146, 152
Metasyrphus corollae, 334
Metopomyza *nigriorbita, 84
Microdon devius, 328; eggeri, 327–8
Mikiola annulipes, 54–5; fagi, 53–6
Mongolomyia, 103, 141–2, 146–7, 152, 156, 158; latitarsis, 146–7
Muscidae, 356
Myathropa florea, 226
Mycetobia, 104
Mycetophilidae, 192
Mydidae, 96, 159, 179, 186
Mythicomyiinae, 95, 110, 156, 158–9
Nematocera, 104, 141
Nemestrinidae, 95–6, 105, 154, 156, 158–9, 179–87, 189
Nemestrininae, 183, 188
Nemestrinoidea, 95–7, 102, 110, 156, 158, 163, 185–90
Nemotelinae, 328
Nemotelus, 328
Neoascia geniculata, 327–8; interrupta, 327; obliqua, 327–8
'Neorhagio', 99, 132
Neorhegmoclemina catharinae, 351–2, 356
Neorhynchocephalus, 179
Neurigona abdominalis, 61–4; pallida, 62, 64; quadrifasciata, 61, 63–4
Nycterimorpha, 179
Nycterimyia, 111, 179; tenestro-inornata, 188
Odontosabula, 99, 186
Olbiogaster, 104
Oligophryne britannica, 161; fungivoroides, 160–1
Oligophryneidae, 160–1
Orthonevra nobilis, 327
Orygma, 79
Ovisargus, 164; gracilis, 166, 168–70, 187
Pachybates, 99–100
Pachyneuridae, 104
Palaeobolbomyia, 131–2, 137; sibirica, 131–2
Palaeobrachyceron, 129–30, 137–8; handlirschi, 129–30; hennigi, 129–30, 137; nagatomii, rohdendorfi, steyskali, 129–30
Palaeophora, 115, 156; ancestrix, 115
Palaeophoridae, 100, 115, 156, 158
Palaeoptiolina, 119–20; 123–6; scobloi, 120–1
Palaeostratiomyia, 129, 132–3, 137; pygmaea, 132
Palembolus, 182
Panopinae, 188
Pantophthalmidae, 96–7, 99–11, 103–4, 110, 141, 143, 156
Paragus tibialis, 327
Parascatopse, 352
Pareremochaetus, 111, 114, 173, 177; minor, 114, 187
Parhadrestia, 100, 139; avata, 139
Parhadrestiinae, 138, 156, 161
Parvisargus malus, 163, 166, 168, 171; peior, 164, 168–9, 171
Passeromyia heterochaeta, 314
Pegomya argyrocephala, bicolor, 264
Pelecorhynchidae, 96, 98–100, 102, 110, 127, 129, 133, 142–3, 145, 147, 152, 185
Pelecorhynchus, 98, 127, 185
Peribaea fissicornis, seticornis, setinervis, 238
Phasia hemiptera, 43
Phoridae, 115, 156, v
Phthiriinae, 136
Platycheirus perpallidus, 328
Platypezidae, 156, xi
Pocota personata, 42
Poecilobothrus muralis, 64; nobilitatus, 62, 64
Pollenia, 334
Polybiomyia macquarti, 223
Probolbomyia, 117, 119, 120–2, 125–6; modesta, 120–2
Procramptonomyiidae, 104
Prohirmoneura jurassica, 181–3, 187
Proptychoptera, 123
Proptychopterina, 123
Prosoeca, 183; florigera, 182
Prosolva, 105–6, 109, 156, 163–4; huabeiensis, 108–9, 166, 169, 171
Prosolva(?) karataviensis, 168–9, 171
Protempididae, 116–7, 155, 158
Protempis antennata, 116–7
Protobrachyceridae, 102, 117, 154, 158
Protobrachyceron, 102, 117; liasinum, 117
Protomphrale, 100, 118, 161; martynovi, 118
Protomphralidae, 100, 118, 141, 156, 161
Protonemestrius, 182–4, 189; bequaerti, handlirschi, longinasus, 183; martynovi, 182–3, 187–8
Protorhagio, 133, 138; capitatus, 133–4
Protovermileo, 117
Pselaphomyia, 328
Pseudoerinna, 99, 110, 127–8, 145, 185
Psilota, 226; anthracina, 39–40, 42, 327; cyanea, 39; decessa, 39–41, 43; victoria, 39, 41
Psychodidae, 356

Pterosargus (s.g.), 165, 166, 170
Ptiolina, 99, 102, 126, 129, 134–5, 137
Ptiolinites, 119, 126, 134, 137–8; cretaceus, 134
Ptychopteridae, 356
Pulicimyia, 233
Puliciphora borinquenensis, 232–4; cavatica, 231–2; glacialis, 233; **kistneri**, 231–3; **nuttingi**, 232–3; occidentalis, 233; omnivora, 233–4; parvula, 232; sobria, 231; suavis, 231, 233; tokyoensis, 233–4; vicinalis, 231–2; virginensis, 233
Rachiceridae, 96, 98, 142, 147, 159
Retinitus, 95
Rhagina, 99
Rhagio, 99, 135–6
Rhagionemestriidae, 162–3, 172, 184–5, 187, 189
Rhagionemestrius, 184–5; rapidus, 184–5, 187
Rhagionempididae, 95, 102, 106, 116–26, 153–7
Rhagionempis, 102, 119–20, 122–6; tabanicornis, 122–3
Rhagionidae, 95–6, 119, 123, 127–38, 141–2, 152, 156, 158–9, 192, 356
Rhagiophryne bianalis, 134–5
Rhaphium appendiculatum, 64
Rhegmoclema, Rhegmoclemina, 351
Rhingia rostrata, 327
Rhinophoridae, xi
Salduba, 140
Sarcophagidae, xi
Scathophaga calida, 77–81; litorea, 77–81; l. ssp. impudica, 81; stercoraria, 77–9, 356
Scathophagidae, 356
Scatopsidae, 104
Scelorhagio, 135–6, 138; mecomastigus, 135
Scenopinidae, 96, 118, 159
Scenopininae, Scenopinus, 118
Schizella, 100
Sciapus loewi, 62–64; platypterus, 61–4; wiedemanni, 62, 64
Sciaridae, 344
Sciomyzidae, 356
screw worm flies, xi
Sepsis punctatum, 334
Sharasargus, 164; ruptus, 169–70
Sharasargus(?) spiniger, 169–70; 187–8
Shevioptera, 118, 123–4; sinitsae, 123–4
Shulmas, 141–2, 147, 152, 158; globiceps, 142, 147
Simuliidae, 104, 344, 356
Sinonemestriidae (fam. nov.), 162–3, 185–6, 189
Sinonemestrius tuanwangensis, 186
Siphona confusa, cristata, geniculata, mesnili, 238
Siphonini, 238
Solomomyia, 100
Solva, 100
Solvidae, 117
Spania, 98–9, 119, 127, 129, 133, 136
Spaniinae, 119
Sphaerophoria, 334
Sphegina, 226; clunipes, 42; verecunda, 327
Stenopteromyia, 179
Stratiomyidae, 95, 98, 100–1, 138–9, 141–2, 148, 156, 159, 161, 328, 356
Stratiomyoidea, 95, 97–8, 100, 102–3, 110, 119, 142, 146, 148, 152, 156
Stratiomyomorpha, 161
Stratiomyopsis, 101, 147–8, 156, 158; robusta, 147–8
Stylospania, 100
Sylvicola, 104
Symphoromyia, 99–100, 134
Sympycnus desoutteri, 64
Syntormon denticulatum, pallipes, 64
Syritta pipiens, 334
Syrphidae, 174, 354, 356
Tabanidae, 96, 98, 159, 356, xi
Tabanoidea, 97–8, 100, 102–4, 119, 141–3, 146, 152, 156, 188
Tachinidae, xi
Thaumaleidae, 356, xi
Themira putris, 344
Therevidae, 96, 159
Thoracochaeta, 79; brachystoma, 81; zosterae, 344
Thripomorpha, 351–2
'Tipula fagi', 53
Tipulidae, 104, 238, xi
Trichacantha, 100
Trichopsideinae, 188
Trichopsomyia flavitarsis, 327
Triglyphus primus, 327
Ussatchovia, 119–20, 122–6; jurassica, 124–5
Vermileonidae, 95, 98–100, 104–6, 111, 117, 119–20, 129, 143
Volucella bombylans, inanis, 292; inflata, 226, 292; pellucens, zonaria, 292
Xanthochlorus ornatus, 62, 64
Xylomya, 100
Xylomyidae, 96, 98, 100, 141–3, 147–8, 156, 159
Xylophagidae, 96–8, 137, 142, 159
Xylophagoidea, 98, 100, 102–4, 110, 129, 142, 146–8, 152, 156, 185–6, 188

Xylophagus, 137
Xylota, 226, 334; florum, tarda, 327
Zarzia, 137; zherichini, 136
Zhangsolva, 101; cupressa, 139–41, 156
Zhangsolvidae (fam. nov.), 95, 101, 139–41, 148, 156, 158, 163

EPHEMEROPTERA

Arthroplea congena, 257
Caenis beskidensis, 94; pseudorivulorum, 94, 263; rivulorum, 94
Ecdyonurus, 260; lateralis group, 257
Electrogena *affinis, 257–63
Ephemeroptera, 192
Heptagenia fuscogrisea, 258, 260–1; lateralis, 257–8, 260–2; longicauda, 257; sulphurea, 258, 260–1

HEMIPTERA

Acrosternum heegeri, 268
Amblytylus delicatus, 265
Aphididae, 44
Aphis abietina, 318
Aploneura lentisci, 76
Aradus cinnamomeus, 265
Arenocoris falleni, 52
Bathysolen nubilis, 52, 265
Berytinus crassipes, minor, 265
Brachycaudus cardui, 85
Cardiastethus fasciiventris, 265
Chilacis typhae, 312
cicadas, 4
Cicadellidae, 192
Cymus glandicolor, 311
Deraeocoris olivaceus, 266
Dicranocephalus agilis, 265
Drepanosiphinae, 44
Drymus brunneus, 311, 313; ryei, 312–3
Elatobium abietinum, 318
Emblethis denticollis, 265
Eremocoris podagricus, 312–3
Eurysa brunnea, douglassi, 307–8; lineata, 307, 309; lurida, 307
Eurysula, 307
Gastrodes grossipes, 311
Gerris paludum, 266
greenflies, 349
Hebrus pusillus, 266
Henestaris halophilus, 265
Heterogaster artemisiae, 265; urticae, 311–3
Hydrometra championana, 65; stagnorum, 65–8
Ischnocoris angustulus, 311
Iziphya bufo, 239–40
Kleidocerys resedae, 312–3; trunculatus, 311
Laingia psammae, 271
Lamproplax picea, 312
Lygus pratensis, 266
Megalonotus chigagra, 312–3; dilatatus, 265, 312; emarginatus, 312–3; e. v. atricornis, 313; praetextatus, 265
Membracidae, 192
Metopoplax ditomoides, 265
Micracanthia marginalis, 266
Microvelia pygmaea, 266
Miridae, 324
Myrmedobia tenella, 265
Neanoecia corni, zirnitsi, 76
Nezara viridula, v.f. smaragdula, v.f. torquata, v.f. viridula, 167–9
Notonecta obliqua, 266
Nysius, 85; ericae, 312; helveticus, 265, 313; thymi, 312
Odontoscelis fuliginosa, lineola, 265
Oliarus panzeri, spp., 309
Ortholomus punctipennis, 313
Palomena prasina, 43
Pemphigidae, 192
Pentatoma metallifera, rufipes, 43
Peritrechus geniculatus, 312–3
Plinthisus brevipennis, 313
Reduviidae, 192
Rhopalus rufus, tigrinus, 265
Rhyparochromus pini, 265
Saldula arenicola, 266
Saltusaphis familiaris, 239
Scolopostethus decoratus, 313; grandis, 312; thomsoni, 311–2
Sehirus luctuosus, 52
Spathocera dahlmanni, 265
Stalia boops, 265
Stictopleurus abutilon, 265
Stygnocoris fuligineus, 313; rusticus, 312; sabulosus, 313
Taphropeltus contractus, 312
Tetraneura ulmi, 76
Thyreocoris scarabaeoides, 52
Trapezonotus arenarius, 312; desertus, 311, 313

HYMENOPTERA

Agelaia ornata, 91; pallipes cuzcoensis, 89
Amonophadnus, 70
Andrena florea, 350; fulvago, humilis, 38; labiata, 350; pilipes, spectabilis, 38
Andrenidae, 350
Anogmus einersbergensis, hohenheimensis, 53
Anoplius caviventris, 350

Anthophorinae, v
ants, 4, 305
Apidae, v, xxi
Aprostocetus elongatus, 56
bees, 3, v
Bethylidae, xxi
Blennocampinae, 68
Caliorini, 69
Caliroa, 192
Chalcididae, 192
Chrysididae, 350, xxi
Chrysis helleni, 350
Colletes fodiens, similis, 334
Colletinae, v
Corpilus, 70
Crabroninae, v
Crossocerus binotatus, 350
Cryptocheilus notatus, 350
Diastrophus rubi, 4
Diodontus insidiosus, 350
Dryinidae, 192
Ectemnius borealis, 350
Encyrtidae, 192
Etroxys Saxeseni, 53
Eulophidae, 192
Eumacepolus grahami, 53–4, 56; saxeseni, 53
Eumenes coarctatus, 350
Eumenidae, 350
Eumeninae, x
Eutomostethus, 70
Euxestonotus, 241–55; **achilles**, 241, 244, 247; acuticornis, 244; **agamemnon**, 241, 252, 256; **ariel**, 241, 248, 251; **astraea**, 241, 244–5, 252; **berolina**, 241, 244, 247; brevicornis, 244; **callisto**, 241–3, 252; **ceres**, 241, 246, 248–9, clavicornis, 246; **deimos**, 241, 248–9; **dione**, 241, 246, 249; **eros**, 241–3; error, 246; flavipes, 246; **ganymedes**, 241, 250, 253; hasselbalchi, 246, 254; **hebe**, 241, 246–7, 252; **hektor**, 241, 246–7; **io**, 241, 250, 253; **japetus**, 241, 245, 247; **juno**, 241, 250, 253; **mimas**, 241, 248, 251; **miranda**, 241, 245, 247; **nestor**, 241, 250, 252–3; nitidus, 250, 254; **oberon**, 241, 246, 249; **pallas**, 241, 250–2; parallelus, 252; **patrokles**, 242, 254–5; **phobos**, 241, 248, 251–2; **photographica**, 241, 248, 251, 254; pini, 246, 254; **priamus**, 241, 252, 254; **rhea**, 241, 244–5; rufidens, 252; **thetys**, 241, 244–5; **titan**, 241–3; **umbriel**, 241, 244–5; **vesta**, 241–3, 254
Fenella, 70
Fenusa, 70
Fenusini, 70
Formica cunicularia, 354
Formicidae, 192, 350, v, xxi
Gorytes bicinctus, 350
Halictus confusus, 238
Haplometrobius, 89, 297 (s.g.)
Heterarthrinae, 69
ichneumonid, 344
Jermakia, 192
Kappa (s.g.), 297
Lasioglossum sexnotatum, zonulus, 334
Lasius brunneus, 350; flavus, 75–6, 354; niger, 75, 354; umbratus, 354
Loopica melanopoda, 69, 71–4; **mizo**, 69, 71–2, 74–5; **nigrescens**, 69, 71–3
Lycaotini, 69
Macropis europaea, 350
Megacanthopus (s.g.), 297
Megachile leachella, 38
Megachilinae, v
Melittidae, 350
Mesopolobus, 53; fagi, 54–6; graminum, juniperinus, 56
Metaneura, 72
Mischocyttarus, 89–93, 293–306; artifex, 293–4, 296–7, 305; artifex-group, 293–306; capichaba, 293; chanchamayoensis, 91; **chloroecus**, 294–5, 297, 300–1; cooperi, 92–3; dimorphus, 293, 295; **filipendulus**, 294–6, 301–3; flavoniger, 301; giffordi, 293; iheringi, 293; iheringi group, 90; illusorius, 293, 295; interruptus, 295, 301, 304–5; **leucoecus**, 293–4, 296, 301–2; **macarenae**, 90–1; **maculipennis**, 295–6, 303; melanopygus, 301; **minifoveatus**, 89, 301; mirificus, 293; **narinensis**, 293–5, 297–8; naumanni, 90, 92–3; nigroclavata, 296–7, 305–6; 296, nigropygidialis, 301; oecothrix, 306; ornatus, 91–2, 300; **polymorphus**, 294, 296, 299–301; peruviensis, 294–5, 297, 301, 304; reflexicollis, 293, 295–6, 306; saussurei, 90; surinamensis group, 89, 298; synoecus, 296–7, 300, 305–6; tertius, 293; thrypticus, 293; vredeni, 90; weyrauchi, 93, 293; ypiranguensis, 293; sp nr mirificus, 303
Mutilla europea, 350
Mutillidae, 350, v, xxi
Myrmica lobicornis, scabrinodis, 354
Nematinae, 69
Nomada pleurostricta, 38
Omalus aeneus, auratus, puncticollis, 52
Omega (s.g.), 297
Omphale lugens, 56

Pararaphidoglossa bicarinata, 300
Paravespula, 279–83; flavipes, 279–81; germanica, 279–80, 282–3; inexpectata, 279; koreensis, 279–81, 283; maculifrons, 279; orbata, 279–81, 283; o. arisana, o. orbata, 279; shidai, structor, 279–80, 282–3; vulgaris, 279–80, 283
Pasteelsia, 69; carinata, constricta, 315, 318; dilatata, 315, 317–8; gracilicornis, 316
Philanthinae, v
Philanthus triangulum, 350
Phymatocera, 70
Phymatoceridia, 69
Phymatoceropsis, 70
Podalonia hirsuta, 38
Polybia, 223
Pompilidae, 350, v, xxi
Psen spooneri, 350
Pteromalus Einersbergensis, 53–4; hohenheimensis, 53
Pygostylus falcatus, multiarticulatus, sticticus, 255
Rhynchium oculatum, 263
Rugovespula (s.g.), 279
Sapyga clavicornis, 350
Sapygidae, 350, xxi
sawflies, 192
Smicromyrme rufipes, 350
Sphecidae, 350, v, xxi
Sphecinae, v
Tenthredinoidea, Tenthredinomorpha, 192
Tetramesa, 56
Tiphia minuta, 350
Tiphiidae, 350, xxi
Tomostethus gracilicornis, 315, 317
Vespa, 279; aegyptica, 45; affinis, 45, 215, 219; analis, 215–22; a. analis, 217; a. barbouri, 215–6; a. eisa, a. insularis-parallela complex, 216; a. kuangsiana, 215–6; a. nagatomii, 216; a. nigrans, 215, 217; a. parallela, 215–7; a. tenebrosa, 216; a. tyrannica, 215, 217–8; arabica, 45–6; basalis, 45; crabro, 306; dybowski, fervida, jurenei, mocsaryana, 45; orbata arisana, 279; orientalis, 45–7; somalica, 45–6; tropica, 45; zavarattii **syn. n.**, 45–6
Vespidae, v, xxi
Vespinae, v
wasps, v

ISOPTERA

Termirostritermes termirostris, 233
termites, xii

LEPIDOPTERA

Brachmia sp., 308
butterflies, 4, 310
caterpillars, 358
Copromorphoidea, 256
Cossus cossus, 343
Epermeniidae, 256
larvae, 206
Lycaenidae, 192, 256
Macroglossum stellatarum, 334
moths, 310
Pieris brassicae, 334
Polygonia c-album, 195, xi
pyralid larvae, 263
Riodinidae, 256
Vanessa atalanta, 334

MECOPTERA

Mecoptera, 192

ODONATA

Platycnemis pennipes nitidula, p. pennipes, 324

ORTHOPTERA

Acrididae, 355
Conocephalus discolor, 60
grasshoppers, 355
Gryllidae, 355
Katydids, 355
locusts, 355
Metrioptera roeselii, 60; r. form diluta, 353
Orthoptera, 60, 192
Pseudomogoplistes squamiger, 230
Stethophyma grossum, 60
Tettigoniidae, 355

PSOCOPTERA

Caecilius macrostigmata, pteridii, 57
Ectopsocus brunneus, hartleyi, 57; pteridii, 57–8
Heterocaecilius monicus, 57–8
Lachiesella pedicularis, 58
Maoripsocus semifuscatus, 57
Paracaecilius lemuris, 57–8
Pentacladus pulchripennis, 57
Peripsocus edwardsi, 57; milleri, 57–8; morulops, 57
Taeniostigma trickettae, 57

THYSANOPTERA

Adiheterothripidae, 345
Aeolothripidae, 345, 347–8
Aeolothrips, 348; ericae, melaleucus, 346–7
Ancothrips, 348
Aulacothrips, 348; dictyotus, 346–7
Cranothrips, 348–9; lauriei, 346–7
Cycadothrips, 348; chadwicki, 346–7
Dactuliothrips, 348; xerophilus, 346–7
Damerothrips gemmatus, 346–8; spp., 346–7
Desmothrips, 348; australis, 346–7; bagnalli, 348
Dorythrips, 348
Echinothrips americanus, 1–3
Erotidothrips spp., 346–7
Erythrothrips, 348
Euceratothrips, 349; marginipennis, 346–7
Fauriella natalensis, 346–7
Fauriellidae, 345, 348
Franklinothrips, 348; vespiformis, 346–7
Heterothripidae, 345, 348
Heterothrips flavicornis, 347
Hindsiothrips bonessi, 319; **navarrensis**, 319–22; pullata, 319, 321; robustisetis, 321
Holarthrothrips tenuicornis, 347
Hoplandrothrips coffeae, flavipes, 346–7
Lenkothrips, 348, sensitivus, 347
Melanthripinae, 348
Melanthrips spp., 346–9
Merothripidae, 345, 347–8
Merothrips, 348; mirus, productus, 346–7
Mymarothrips, 349; ritchianus, 346–7
Oligothrips spp., 347
Orothrips, 348; priesneri, 346–7
Panchaetothripinae, 3
Phlaeothripidae, 345
Rhipidothrips, 348; cahirensis, 346–7
Sericothripini, 3
Thripidae, 345
Thripinae, 2
Thripini, 3
Thysanoptera, 324
Tylothrips osborni, 319, 322–3
Uzelothripidae, 345

TRICHOPTERA

Trichoptera, 192

GROUPS OTHER THAN INSECTA AND ARACHNIDA

Aves, 81, 205, 314
Bacteria, 228
Coelenterata, 81
Mammalia, 344
Myriapoda, 266
Protozoa, 229

ERRATA AND CORRIGENDA

Page	Line	For	Read
95	19 (up)	questionally	questionably
103	1 (up)	Krezemiński	Krzemiński
256	3	DEVRIES	DE VRIES
xxi	6	*Cyuphea*	*Cyphea*

Printed in England by Attwood Dawes Printers, K